

Methicillin-resistant *Staphylococcus aureus* (MRSA) Update, Q3 of 2017/18

March 2018

Summary Table

	Q3 2017/18	Previous quarter (Q2 2017/18)	Same quarter of previous year (Q3 2016/17)	Year-to- date 2017/18
Total new MRSA cases identified	762	756	716	2,134
Number of new MRSA cases associated with the reporting facility	331	359	357	969
Total inpatient days	726,096	690,507	715,803	2,060,870
Rate of MRSA associated with the reporting facility per 10,000 inpatient days (95% CI)	4.6 (4.1-5.1)	5.2 (4.7-5.8)	5.0 (4.5-5.5)	4.7 (4.4-5.0)

Highlights

- The provincial rate of new MRSA cases associated with the reporting facility in Q3 2017/18 was 4.6 per 10,000 inpatient days, which was lower than the previous quarter (5.2 per 10,000 inpatient days). However, the difference was not statistically significant.
- The MRSA rate in Q3 of 2017/18 was not significantly different from the same quarter of the previous year (5.0 per 10,000 inpatient days in Q3 of 2016/17).
- The provincial rate of MRSA has not changed significantly from Q1 of 2013/2014 to Q3 of 2017/2018.

What is Methicillin-resistant *Staphylococcus aureus* (MRSA)?

MRSA is a type of *Staph* bacteria that has become resistant to many antibiotics, including methicillin, penicillin, and amoxicillin, and is thus more difficult to treat. MRSA often lives on the skin or in the nose of healthy people without causing symptoms (this is called colonization). It can, however, cause skin and other infections. Most infections are minor, such as pimples and boils. Serious infections — such as severe wound infections, pneumonia, or septicaemia (infection in the bloodstream) — can result in life-threatening illness or, if left untreated, death. Those with weakened immune systems and chronic illnesses are more susceptible to developing an infection.

How is MRSA transmitted?

MRSA is primarily spread by skin-to-skin contact or through contact with surfaces contaminated with the bacteria. About 2 in 100 people in the community carry MRSA. People who carry MRSA but do not have signs of infection can spread the bacteria to others. It has shown to spread easily in healthcare settings; therefore hospital patients and residents in residential care facilities are at a higher risk of acquiring MRSA.

How to prevent MRSA transmission?

Clean your hands often with soap and water or use an alcohol-based hand rub (sanitizer); do not share personal items; cover your wounds or cuts with clean, dry bandages until healed; and wash used sheets, towels, and clothes with warm water and laundry detergent. The spread of MRSA can be prevented in healthcare settings through adherence to infection control measures such as contact precautions, careful cleaning of patient rooms and medical equipment, and good hand hygiene on the part of physicians, nursing staff, and other healthcare providers.

Why is MRSA being monitored in BC hospitals?

MRSA bacteria have many virulence factors that enable them to cause disease. For example, MRSA is a cause of healthcare-associated bloodstream and catheter-related infections. MRSA is also a common cause of community-associated infections, especially skin and soft tissue infections, and can also cause necrotizing pneumonia. Active surveillance (e.g., screening of high-risk individuals) can identify patients colonized with MRSA so that precautions can be taken to prevent transmission to other patients.

Where to find information about MRSA

The PICNet website (www.picnet.ca) provides provincial guidance, toolkits, and related resources for the prevention and control of MRSA and other antimicrobial organisms, as well as the surveillance protocol and reports on MRSA in BC. If you have questions or suspect that you have MRSA, please contact your doctor or healthcare provider.

This quarterly update presents the latest data on the incidence and trends of healthcare-associated (HCA) cases of MRA that were newly identified among inpatients in the last five years. In the following graphs,

- 1) MRSA cases that were associated with another healthcare facility, community-associated, or of unknown origin were not included.
- 2) The data were aggregated by fiscal quarter for each health authority except Provincial Health Services Authority (PHSA), which aggregated the data by calendar quarter. The time frame of each fiscal quarter varied by fiscal year and there were more days in the fourth quarter (Q4) than in other three quarters (Q1, Q2, and Q3) of each fiscal year.
- 3) The line in each graph represents the overall linear trend over time.
- 4) Direct comparison of the number of cases or the rate between health authorities is not recommended due to variations in case finding strategies and the application of MRSA case classification for surveillance among the health authorities.

Figure 1. Provincial rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and quarter, 2013/14 - 2017/18, British Columbia




Figure 2. Rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and quarter, 2013/14 - 2017/18, Interior Health




Figure 3. Rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and quarter, 2013/14 - 2017/18, Fraser Health




Figure 4. Rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and quarter, 2013/14 - 2017/18, Vancouver Coastal Health¹





Figure 5. Rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and quarter, 2013/14 - 2017/18, Island Health²



¹ Includes acute care facilities of Providence Health Care (PHC)

² The data include two new hospitals opened during Q3 of 2017/18 and historical data from two closed hospitals.

Figure 6. Rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and quarter, 2013/14 - 2017/18, Northern Health





Figure 7. Rate and number of new cases of MRSA associated with the reporting facility, by fiscal year and calendar quarter, 2013/14 - 2017/18, Provincial Health Services Authority



Provincial Infection Control Network of BC (PICNet)
1001 West Broadway, Suite 504
Vancouver, BC V6H 4B1
Tel: 604-875-4844 x 22985
Fax: 604-875-4373
Website: www.picnet.ca
Email: picnet@phsa.ca

Disclaimer

The purpose of this report is to provide information to healthcare providers, decision-makers, patients, and the public on healthcare-associated infections identified among the patients admitted to acute care facilities. This report may be used, in whole or in part, as the basis for infection prevention and control practices to improve the quality of healthcare services. PICNet does not warrant or assume any legal liability or responsibility for the accuracy, completeness, or usefulness of any information in the report; neither does it intend to provide specific medical advice. Commercial uses are prohibited without express written permission.